Appl. No.: 10/659,475

Response dated February 16, 2006

Reply to Notice of Appeal filed December 12, 2005

Amendments to the Claims

The listing of claims replaces all prior versions, and listings, of claims in the application:

Listing of Claims:

- 1. (Currently amended) A carotenoid composition comprising lycopene suspended or dispersed in a medium-chain triglyceride, wherein the medium-chain triglyceride is derived from esterification of a substantially pure medium chain fatty acid and substantially pure glycerol which maintains the concentration of lycopene for at least three months at 25°C.
- (Cancelled).
- 3. (Previously presented) A composition according to claim 1, wherein the fatty acid is a C6 to C12 fatty acid.
- 4. (Previously presented) A composition according to claim 1, wherein the fatty acid is a C8 to C10 fatty acid.
- (Previously presented) A composition according to claim 1, wherein the fatty acid is selected from the group consisting of caprylic acid, capric acid, lauric acid, and mixtures thereof.
- 6. (Previously presented) A composition according to claim 1, comprising up to 10% by weight lycopene.
- 7. (Previously presented) A composition according to claim 6, comprising between 4%

Appl. No.: 10/659,475
Response dated February 16, 2006
Reply to Notice of Appeal filed December 12, 2005

and 5% by weight lycopene.

- 8. (Previously presented) A composition according to claim 6, comprising up to 98% by weight medium chain triglyceride.
- 9. (Previously presented) A composition according to claim 8, comprising between 20% and 80% by weight medium chain triglyceride.
- 10. (Previously presented) A composition according to claim 9, comprising between 30% and 70% by weight medium chain triglyceride.
- 11. (Previously presented) A composition according to claim 8, further comprising an oil-soluble antioxidant.
- 12. (Previously presented) A composition according to claim 11, wherein the oil-soluble anti-oxidant is selected from the group consisting of tocopherols, butylated hydroxy anisole, butylated hydroxytoluene, propyl gallate, ethoxyquin and ascorbyl palmitate, natural tocopherols, and mixtures thereof.
- 13. (Previously presented) A composition according to claim 12, wherein the tocopherol is selected from the group consisting of beta, gamma-, and delta-tocopherols and mixtures thereof.
- 14. (Previously presented) A composition according to claim 11, wherein the antioxidant comprises up to 15% of the composition.

Appl. No.: 10/659,475
Response dated February 16, 2006
Reply to Notice of Appeal filed December 12, 2005

- 15. (Previously presented) A composition according to claim 14, wherein the antioxidant comprises between 5% and 10% by weight of the formulation.
- 16. (Currently amended) A method of providing a bioavailable antioxidant comprising orally administering to a host an effective amount of lycopene suspended or dispersed in a medium chain triglyceride, wherein the medium chain triglyceride is derived from esterification of a substantially pure medium chain fatty acid and substantially pure glycerol which maintains the concentration of lycopene for at least three months at 25°C.